



Contribution ID: 38

Type: **Presentation**

Progress towards barium daughter tagging in Xe^{136} decay using single molecule fluorescence imaging

Sunday, 24 September 2017 11:40 (15 minutes)

The existence of Majorana fermions is of great interest as it may be related to the asymmetry between matter and anti-matter particles in the universe. However, the search for them has proven to be a difficult one. Neutrino-less Double Beta decay (NLDB) offers a possible opportunity for direct observation of a Majorana Fermion. The rate for NLDB decay may be as low as $\approx 1 \text{ count/ton/year}$ if the mass ordering is inverted. Current detector technologies have background rates between 4 to 300 $\text{count/ton/year/ROI}$ at the 100kg scale which is much larger than the universal goal of 0.1 $\text{count/ton/year/ROI}$ desired for ton-scale detectors. The premise of my research is to develop new detector technologies that will allow for a background-free experiment. My current work is to develop a sensor that will tag the daughter ion Ba^{++} from the Xe^{136} decay. The development of a sensor that is sensitive to single barium ion detection based on the single molecule fluorescence imaging technique is the major focus of this work. If successful, this could provide a path to a background-free experiment.

Primary author: MCDONALD, Austin (UTA)

Presenter: MCDONALD, Austin (UTA)

Session Classification: Sunday Morning 2

Track Classification: Detector techniques (HV, cryogenics, purification, calibration, etc.)